





## Conclusion

Fasting in patients with CVT using OCPs causes significant increase in focal neurological deficit and hemorrhagic infarction at presentation, which also increases the hospital stay. However, it does not significantly change long-term prognosis and mortality of CVT in women using OCPs.

## Conflict of Interests

The authors declare no conflict of interest in this study.

## Acknowledgments

We thank all the patients and their families for their cooperation.

## References

1. Stam J. Thrombosis of the cerebral veins and sinuses. *N Engl J Med* 2005; 352(17): 1791-8.
2. Amoozegar F, Ronksley PE, Sauve R, Menon BK. Hormonal contraceptives and cerebral venous thrombosis risk: A systematic review and meta-analysis. *Front Neurol* 2015; 6: 7.
3. Borhani HA, Ashjazadeh N, Safari A, Cruz-Flores S. Cerebral venous sinus thrombosis in Iran: Cumulative data, shortcomings and future directions. *Iran Red Crescent Med J* 2012; 14(12): 805-10.
4. Saeidi M, Foroughipour M, Sasannezhad P, Melat Ardakani A, Azarpazhouh MR. The relation between short course oral contraceptive consumption and cerebral vein thrombosis in Ramadan. *Iran J Neurol* 2008; 7(23): 260-5.
5. Azin H, Ashjazadeh N. Cerebral venous sinus thrombosis--clinical features, predisposing and prognostic factors. *Acta Neurol Taiwan* 2008; 17(2): 82-7.
6. Coutinho JM, Ferro JM, Canhao P, Barinagarrementeria F, Cantu C, Boussier MG, et al. Cerebral venous and sinus thrombosis in women. *Stroke* 2009; 40(7): 2356-61.

Proof Version