

Effects of hydroalcoholic extract of *Coriandrum sativum* on oxidative damage in pentylenetetrazole-induced seizures in rats

Received: 12 Dec 2014
Accepted: 16 Jan 2015

Reza Karami¹, Mahmoud Hosseini², Toktam Mohammadpour³, Ahmad Ghorbani⁴, Hamid Reza Sadeghnia⁴, Hassan Rakhshandeh⁴, Farzaneh Vafaei³, Mahdi Esmaeilizadeh⁵

¹ Student Research Committee, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

² Neurocognitive Research Center AND Department of Physiology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

³ Neurogenic Inflammation Research Center AND Department of Physiology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

⁴ Pharmacological Research Center of Medicinal Plants AND Department of Pharmacology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

⁵ Esfarayen Faculty of Medical Sciences, Esfarayen, Iran

Keywords

Coriandrum sativum, Pentylenetetrazole, Seizures, Rat, Oxidative Stress, Brain

Abstract

Background: An important role for oxidative stress, as a consequence of epileptic seizures, has been suggested. *Coriandrum sativum* has been shown that have antioxidant effects. Central nervous system depressant effects of *C. sativum* have also been reported. In this study, the effects of hydroalcoholic extract of aerial parts of the plants on brain tissues oxidative damages following seizures induced by pentylenetetrazole (PTZ) was investigated in rats.

Methods: The rats were divided into five groups and treated: (1) Control (saline), (2) PTZ (90 mg/kg, i.p.), (3-5) three doses (100, 500 and 1000 mg/kg of *C. sativum* extract (CSE) before PTZ. Latencies to the first minimal clonic seizures (MCS) and the first generalized tonic-clonic seizures (GTCS) were recorded. The cortical and hippocampal tissues were then removed for biochemical measurements.

Results: The extract significantly increased the MCS and GTCS latencies ($P < 0.01$, $P < 0.001$) following

PTZ-induced seizures. The malondialdehyde (MDA) levels in both cortical and hippocampal tissues of PTZ group were significantly higher than those of the control animals ($P < 0.001$). Pretreatment with the extract prevented elevation of the MDA levels ($P < 0.010$ – $P < 0.001$). Following PTZ administration, a significant reduction in total thiol groups was observed in both cortical and hippocampal tissues ($P < 0.050$). Pretreatment with the 500 mg/kg of the extract caused a significant prevention of decreased in total thiol concentration in the cortical tissues ($P < 0.010$).

Conclusion: The present study showed that the hydroalcoholic extract of the aerial parts of *C. sativum* possess significant antioxidant and anticonvulsant activities.

Introduction

Epilepsy is a common neurological disease, which affects approximately 1% of the population.¹ It is characterized by abnormal episodic bursts of electrical activity in neurons, which is followed by a significant impact on the behavior of the affected patients.² An important role for oxidative stress both as a consequence and as a cause of epileptic seizures has been suggested.³ It has been reported that production

of free radicals increases during seizures, which is lead to oxidative damage to lipids, DNA and susceptible proteins.⁴

Due to high levels of membrane lipid constituents, the central nervous system (CNS) is very susceptible to oxidative injury.⁵ In addition, oxidative damage plays a significant role in the pathogenesis of various CNS disorders and neurobehavioral impairments.⁵ The functional impairments of CNS, which occur during seizures have also been suggested to be at least in part, related to the brain tissues oxidative damages.⁶ Furthermore, the anticonvulsant activities of several agents with antioxidant effects such as melatonin, vineatrol, trans-resveratrol and alpha lipoic acid have been documented.^{7,8} There are also some reports that reactive oxygen species (ROS) may underlie the convulsant and neurotoxic effects of pentylenetetrazole (PTZ).⁹ The results of human and animal studies imply that epilepsy and seizures are lead to the brain tissues oxidative damages, especially in the cortical and hippocampal regions, which are accompanied with cognitive, learning and memory deficits.^{4,6,8,10}

Medicinal plants are good sources to find new therapeutic agents for human diseases. Coriandrum sativum, an annual herb belonging to the Apiaceae family, has been reported to have a wide range of biological activities including sedative-hypnotic, antidiabetic, hypolipidemic, and hepatoprotective effects.¹¹⁻¹⁵ Experimental studies have also revealed a strong antioxidant activity for *C. sativum* that is superior to the well-known antioxidant agents like ascorbic acid.¹⁵⁻²⁰ In our previous work, we found that the hydroalcoholic extract of aerial parts of this plant bearing some compounds with the hypnotic effects.²¹ Regarding the antioxidant and CNS depressant effects of *C. sativum*, we aimed to evaluate the possible protective effects of aerial parts of the plant on PTZ-seizures and the brain tissues oxidative damages in rats.

Materials and Methods

PTZ was purchased from Sigma-Aldrich Company (St. Louis, USA). Other chemical compounds such as thiobarbituric acid (TBA), trichloroacetic acid (TCA), hydrochloric acid (HCL), ethylenediaminetetraacetic acid (EDTA) and 2, 2'-dinitro-5, 5'-dithiodibenzoic acid (DTNB) were bought from Merck Company.

In this study, 40 virgin male Wistar rats, 250 ± 20 g in weight were used. The animals were maintained at the animal house under controlled conditions including 12 h light and dark cycle, 22-24 °C temperature and appropriate humidity with laboratory chow and water provided ad libitum.

The animals were randomly divided into five groups and treated ($n = 8$ in each group) as follows:

(1) Control (saline), (2) PTZ, (3) *C. sativum* extract (CSE) 100 mg/kg (CSE 100) + PTZ, (4) CSE 500 mg/kg (CSE 500) + PTZ and (5) CSE 1000 mg/kg (CSE 1000) + PTZ. The doses were chosen regarding our previous study.²¹ The number of animals was also based on our previous studies.^{9,21-24}

The animals in groups 2-5 were treated intraperitoneally (i.p.) with saline or the extract 30 min before i.p. injection of a single dose (90 mg/kg) of PTZ. In our previous works, we showed that PTZ in this dose induces generalized tonic-clonic seizures (GTCS) in rats.^{9,22,24,25} The time interval between injection of the extract and PTZ was chosen regarding our previous work in which injection of the extract 30 min before injection of pentobarbital increased the sleeping time.²¹

The cortical and hippocampal regions were then removed for biochemical measurements. In the control group, saline was injected instead of both PTZ and CSE and the brain tissues were removed without inducing the seizures. All efforts were made to maintain the animals in good general health, in accordance with the European Communities Council Directive (2010/63/UE). All behavioral tests were conducted between 10:00 and 14:00 O'clock. Animal handling and all related procedures were confirmed by the Mashhad University of Medical Sciences, Iran, Ethical Committee.

The aerial parts (leaves, stems, twigs) of *C. sativum* were collected from Neyshabur, Iran. The identity of the plant was confirmed and for future reference a voucher specimen (10068) was deposited at the herbarium of School of Pharmacy (Mashhad University of Medical Sciences). To prepare the hydroalcoholic extract, the plant materials (50 g) were dried and extracted with 300 ml ethanol-water (70/30, v/v) using a Soxhlet apparatus. The extract reduced to dryness with a rotary vacuum evaporator (Stuart RE300, UK).²⁶

In order to observe ictal behavior, PTZ was injected and the animals were placed in a Plexiglas arena (30 cm × 30 cm × 30 cm) on the day of the experiment. The animals were observed during 60 min after PTZ (90 mg/ kg) administration.^{9,22,24,25,27} The behavioral responses of the animals to PTZ administration were evaluated using these criteria: latency to the first minimal clonic seizure (MCS), incidence of MCS, latency to the first GTCS, incidence of GTCS, protection percentage against GTCS and protection percentage against mortality.²³⁻²⁵

After behavioral study, the rats were quickly decapitated under deep sodium pentobarbital anesthesia, their brains were removed and the cortical and hippocampal regions were separated and conserved for biochemical measurements. The animals were killed by a competent person with a

minimum pain, suffering, and distress. The method was performed as set out in that Annex IV of the guidelines from Directive EU/2010/63 of the European Parliament.

For total thiol (SH) content measurement, the cortical and hippocampal regions were dissected on an ice-cold surface and homogenized in iced-cold phosphate-buffered saline to give 10% homogeny. Total SH groups were measured using DTNB as the reagent. This reagent reacts with the thiol groups to produce a yellow colored complex, which has a peak absorbance at 412 nm. Briefly, 1 ml Tris-EDTA buffer (pH = 8.6) was added to 50 µl of the brain homogenates, and the sample absorbance was read at 412 nm against Tris-EDTA buffer alone (A₁). Then, 20 µl DTNB reagents (10 mm in methanol) were added to the mixture and after 15 min (stored in laboratory temperature), the sample absorbance was read again (A₂). The absorbance of DTNB reagent was also read as a blank (B). Total thiol concentration (mm) was calculated from the following equation.^{9,28-30}

Total thiol concentration (mM) = (A₂-A₁-B) × 1.07/0.05 × 13.6

Malondialdehyde (MDA) levels, as an index of lipid peroxidation, were also measured. MDA reacts with TBA as a TBA reactive substance to produce a red colored complex, which has a peak absorbance at 535 nm. The TBA/TCA/HCL reagent was added to the homogenates, and the solution was heated in a water bath for 40 min. After cooling, the whole solutions were centrifuged within 1000 g for 10 min. The absorbance was measured at 535 nm.^{9,28-30} The MDA concentration was calculated as follows: C (M) = Absorbance/(1.56 × 10⁵)

All data were expressed as mean ± standard error of the mean and analyzed by using ANOVA, followed by Tukey's post-hoc comparison test. P < 0.0500 were considered to be statistically significant.

Results

Effect of *C. sativum* on PTZ-induced seizures

All the animals in different groups (except the control group, which did not receive PTZ) showed MCS and GTCS following administration of a high dose of PTZ. Data analysis using one-way ANOVA showed that there was a significant difference between the groups in MCS latencies (F_{3,28} = 19.65, P < 0.0001). MCS latencies in the extract pre-treated groups were significantly higher than that of PTZ group. When compared with PTZ group (61.66 ± 4.76 s), 100, 500 and 1000 mg/kg of the extract significantly (P < 0.0100 to P < 0.0010) increased the MCS latencies to 77.3 ± 2.07, 93.5 ± 4.35 and 339.3 ± 58.96 s, respectively (Figure 1).

Data analysis using one-way ANOVA also showed that there was a significant difference between the

groups in GTCS latencies (F_{3,28} = 24.13, P < 0.0001). The GTCS latencies in the animals, which had received 100, 500 and 1000 mg/kg of CSE before PTZ, were 183 ± 4.69, 341.88 ± 44.16, and 710.3 ± 98.84 c, respectively. All 3 doses of the extract significantly increased the GTCS latencies (P < 0.0100 to P < 0.0010) compared with PTZ group (114 ± 1.8 c) (Figure 2). There were no significant differences in mortality rate following PTZ administration between groups.




Figure 1. Latencies to minimal clonic seizures (MCS) onsets in pentylenetetrazole (PTZ), *C. sativum* extract (CSE) 100 mg/kg (CSE 100)-PTZ, CSE 500 mg/kg (CSE 500)-PTZ, CSE 1000 mg/kg (CSE 1000)-PTZ groups. The animals were treated with saline or CSE (100, 500 or 1000 mg/kg) before a single injection (90 mg/kg) of PTZ; ** P < 0.010; *** P < 0.001 as compared to PTZ group




Figure 2. Latencies to generalized tonic-clonic seizures (GTCS) onsets in pentylenetetrazole (PTZ), *C. sativum* extract (CSE) 100 mg/kg (CSE 100)-PTZ, CSE 500 mg/kg (CSE 500)-PTZ, CSE 1000 mg/kg (CSE 1000)-PTZ groups. The animals were treated with saline or CSE (100, 500 or 1000 mg/kg) before a single injection (90 mg/kg) of PTZ; ** P < 0.010; *** P < 0.001 as compared to PTZ group

Effect of *C. sativum* on brain tissues oxidative damage

Data analysis using one-way ANOVA showed that there was a significant difference between the groups in MDA concentrations of cortical tissues ($F_{4,35} = 6.93$, $P < 0.001$). The MDA levels in cortical regions of PTZ group were significantly higher than those of control animals ($P < 0.001$) (Figure 3). As shown in figure 3, pretreatment with both 100 and 1000 mg/kg of the extract resulted in a significant reduction in the free radical-mediated lipid peroxidation as indicated by a decrease in the MDA levels ($P < 0.001$ and $P < 0.010$, respectively).

Data analysis using one-way ANOVA also showed that there was a significant difference between the groups in total thiol contents of cortical tissues ($F_{4,35} = 6.78$, $P < 0.001$). Following PTZ administration, a significant reduction in total SH groups in cortical samples was observed ($P < 0.050$, Figure 4). Pretreatment with 500 mg/kg of the extract prevented of decreased total thiol concentration in cortical tissues, as compared with PTZ group ($P < 0.010$).




Figure 3. Comparison of the malondialdehyde (MDA) levels in cortical tissues of control, pentylenetetrazole (PTZ), *C. sativum* extract (CSE) 100 mg/kg (CSE 100)-PTZ, CSE 500 mg/kg (CSE 500)-PTZ, CSE 1000 mg/kg (CSE 1000)-PTZ groups. The animals were treated with saline or CSE (100, 500 or 1000 mg/kg) before a single injection (90 mg/kg) of PTZ; The animals in control group received saline instead of PTZ; *** $P < 0.001$ as compared to control group; ++ $P < 0.010$; +++ $P < 0.001$ as compared to PTZ group

Data analysis using one-way ANOVA showed that there was a significant difference between the groups in MDA concentrations of hippocampal tissues ($F_{4,35} = 24.53$, $P < 0.0001$). The MDA levels in the hippocampal regions of PTZ group were significantly higher than those of control animals ($P < 0.001$) (Figure 5). The results also showed that all three doses of CSE prevented the elevation of MDA concentration in hippocampal tissues ($P < 0.001$ for all, Figure 5).

Data analysis using one-way ANOVA also showed that there was a significant difference between the

groups in total thiol contents of hippocampal tissues ($F_{4,35} = 3.19$, $P < 0.050$). Following PTZ administration, a significant reduction in total SH groups in the hippocampal samples was observed ($P < 0.050$, Figure 6). There were no significant differences between CSE treated rats and PTZ group when total thiol content in hippocampal tissues was compared (Figure 6).




Figure 4. Comparison of the total SH groups in cortical tissues of control, pentylenetetrazole (PTZ), *C. sativum* extract (CSE) 100 mg/kg (CSE 100)-PTZ, CSE 500 mg/kg (CSE 500)-PTZ, CSE 1000 mg/kg (CSE 1000)-PTZ groups. The animals were treated with saline or CSE (100, 500 or 1000 mg/kg) before a single injection (90 mg/kg) of PTZ. The animals in the control group received saline instead of PTZ; * $P < 0.0500$ as compared to Control group; ++ $P < 0.0100$ as compared to PTZ group




Figure 5. Comparison of the malondialdehyde (MDA) levels in hippocampal tissues of control, pentylenetetrazole (PTZ), *C. sativum* extract (CSE) 100 mg/kg (CSE 100)-PTZ, CSE 500 mg/kg (CSE 500)-PTZ, CSE 1000 mg/kg (CSE 1000)-PTZ groups. The animals were treated with saline or CSE (100, 500 or 1000 mg/kg) before a single injection (90 mg/kg) of PTZ. The animals in the control group received saline instead of PTZ; *** $P < 0.0010$ as compared to control group; +++ $P < 0.001$ as compared to PTZ group




Figure 6. Comparison of the total SH groups in hippocampal tissues of control, pentylenetetrazole (PTZ), *C. sativum* extract (CSE) 100 mg/kg (CSE 100)-PTZ, CSE 500 mg/kg (CSE 500)-PTZ, CSE 1000 mg/kg (CSE 1000)-PTZ groups. The animals were treated with saline or CSE (100, 500 or 1000 mg/kg) before a single injection (90 mg/kg) of PTZ; The animals in control group received saline instead of PTZ; * $P < 0.050$ as compared to control group.

Discussion

Oxidative stress is a basis etiology for many neurological and neurodegenerative disorders. Previous studies demonstrated that oxidative stress plays an important role in the pathogenesis of epileptic seizures.^{3,4,31} Elevation of free radical levels during seizures^{7,31} has been well documented therefore, it is suggested that oxidative stress has an important role in the brain damages due to epilepsy.^{9,32} Furthermore, the brain tissues oxidative damage contributes to the complications of seizures and epilepsy including cognitive, learning and memory impairments.^{2,10} In the present study the possible protective effects of *C. sativum* aerial parts on PTZ-induced seizures and the brain tissues oxidative damages was investigated. PTZ, is a selective inhibitor of the chloride channel which is coupled to the gamma-aminobutyric acid receptor.³³ It is a well-known chemoconvulsant which is frequently used for evaluation of antiepileptic drugs.^{34,35} A high dose of PTZ induces a continued seizure activity which progress from mild myoclonic jerks to face and forelimbs clonus without loss of righting reflex (which is known as MCS), to clonic seizures of limbs with loss of righting reflex which is followed by full tonic extension of both forelimbs and hindlimbs (GTCS).³⁶ PTZ has been repeatedly used in 90-100 mg/kg to induce MCS and GTCS seizures.^{9,22-25,27,28} We also previously showed that PTZ-induced seizures are accompanied with brain tissues oxidative damage.⁹ The contribution of ROS to the neurotoxic effects of

PTZ has also been suggested.^{37,38} Similarly, in the present study, we observed an increase in MDA levels and a reduction in total SH groups in the brains of the animals subjected to PTZ-induced seizure. Increase in ROS production, including superoxide anions, hydroxyl radicals, and hydrogen peroxide, in the brains subjected to seizures, have been well documented.^{39,40} It has been suggested that oxidative damages of brain tissues by free radicals may lead to psychiatric or cognitive problems such as depression, anxiety and memory loss.^{31,41} The decreasing in the life span, which has been reported in the epileptic persons may also be at least in part, due to oxidative damages.⁴² Oxidative stress has also been suggested as a link between aging and seizure.⁴³ In the present study, we assessed the effect of the extract by studying its effect on lipid peroxidation, which was measured in terms of MDA concentrations. Studies with animal models using the MDA assay have generally reported an increased lipid peroxidation in the brain tissues in seizures and epilepsy.^{37,44} In our experiments, we observed a significant increase of lipid peroxidation in the both hippocampal and cortical tissues which was prevented by 100 and 1000 mg/kg of the extract however, 500 mg/kg of the extract prevented MDA elevation in hippocampal but not cortical tissues. These results are in consistent with the recently reported protective effects of the plant against hippocampal tissues oxidative damage.⁴⁵⁻⁴⁷ It has been previously suggested the brain regions are differently vulnerable to increased lipid peroxidation. For example Shila et al. showed that lipid peroxidation was increased to the highest level in hippocampus among the brain regions, followed by cortex in arsenic intoxicated rats which was attributed to the different levels of iron contents.⁴⁸ Regarding the results of the present study, it seems that the plant extract was more effective to prevent of increased lipid peroxidation due to PTZ-induced seizures in hippocampal tissues compared to cortical tissues. In consistent with these results, Velaga et al. also reported that the hydroalcoholic extract of *C. sativum* seeds was more effective to decrease lipid peroxidation in hippocampus of lead intoxicated rats, compared to the cortical, cerebellum and brain stem tissues.⁴⁵ It has also been reported that training exercise and vitamin E were more effective to reduce lipid peroxidation in the hippocampal tissues of aged rats compared with the cortical tissues.⁴⁹ SH groups are also known to be sensitive to oxidative damage and depleted following an oxidative insult⁵⁰ therefore, we studied the effect of the extract on total thiol concentrations in the brain tissues after seizures. Similar to other studies, thiol groups were decreased in the brain following seizures injury. As mentioned

earlier, CSE prevented PTZ-induced thiol depletion only in the cortex and not hippocampus. One possible explanation for this observation might be due to brain regional-dependent glutathione (GSH) metabolism, as a major source of thiol group.⁵¹ It has been observed that brain GSH concentrations varied in the range of 0.2-10 mM.⁵² It was reported that GSH levels are highest in cortex, followed by hippocampus and brain stem.^{53,54} Consistent with this finding, some studies reported a strong antioxidant activity for *C. sativum*.¹⁵⁻²⁰ Furthermore, we showed that this activity of *C. sativum* was accompanied by an anticonvulsant effect as it increased both MCS and GTCS latencies. In keeping with these observations, the anticonvulsant activity of several agents with antioxidant effect such as melatonin, vineatrol, trans-resveratrol and alpha lipoic acid have also been shown.^{7,8} Using pentobarbital-induced hypnosis animal model, we previously found that the injection of 50, 100, 200 and 400 mg/kg of hydroalcoholic extract of *C. sativum* before pentobarbital increased the sleep time of the rats.¹² Even a higher dose of the extract has also been previously used in the rats without observation of any toxic effect.⁵⁵ In the current study, therefore we used comparable doses of CSE 30 min before PTZ injection to test its anticonvulsive effect. Previously, Hosseinzadeh and Madanifard reported an anticonvulsant effect for the aqueous and ethanolic extracts of *C. sativum* seeds.⁵⁶ The same effect was found by Emamghoreishi and Heidari-Hamedani for aqueous and hydroalcoholic extracts and essential oil of the seeds.⁵⁷ Regarding the results of present study the beneficial effects of aerial parts of the plant on seizures and its complications such as brain tissues oxidative is suggested however, further studies using other animal models are needed to be done in the future. The electrophysiological studies are also suggested for future.

However, no pharmacological studies have been yet evaluated the anticonvulsant activity of aerial parts of this plant. These parts of *C. sativum* are widely consumed as a vegetable all over the world. With the present study, we showed that aerial parts of *C. sativum* are effective in protection against seizures

and oxidative stress induced by PTZ.

In the present study, the chemical compound(s) responsible for these effects of *C. sativum* were not identified and needs to be more investigated in future studies. The presence of the flavonoids such as quercetin has been reported in CSE.⁵⁸ On the other hand, it has been shown that the flavonoids have considerable anticonvulsant effects.^{59,60} Regarding sedative and CNS depressant effects of flavonoids such as quercetin, these effects could be attributed to the affinity of the compounds for the central benzodiazepine receptors.⁶¹⁻⁶⁵ The beneficial effect of linalool in PTZ as well glutamate-related seizure models has been suggested.^{66,67} It might be suggested the beneficial effects of the extract which was observed in the present study, might be at least in part, due to linalool which is one the main constituents of coriander and has considerable antioxidant effects.⁶⁸

Conclusion

The present data demonstrate that the hydroalcoholic extract of *C. sativum* aerial parts possesses anticonvulsant activity. This activity is accompanied by an antioxidant effect in brain tissues. Isolation of the active compound(s) from the extract needs to be done in the future.

Conflict of Interests

The authors declare no conflict of interest in this study.

Acknowledgments

The authors would like to thank the Vice Presidency of Research, Mashhad University of Medical Sciences, for its financial supports.

How to cite this article: Karami R, Hosseini M, Mohammadpour T, Ghorbani A, Sadeghnia HR, Rakhshandeh H, et al. Effects of hydroalcoholic extract of *Coriandrum sativum* on oxidative damage in pentylenetetrazole-induced seizures in rats. *Iran J Neurol* 2015; 14(2): 59-66.

References

1. Sander JW. The epidemiology of epilepsy revisited. *Curr Opin Neurol* 2003; 16(2): 165-70.
2. Meador KJ. Cognitive outcomes and predictive factors in epilepsy. *Neurology* 2002; 58(8 Suppl 5): S21-S26.
3. Patel M. Mitochondrial dysfunction and oxidative stress: cause and consequence of epileptic seizures. *Free Radic Biol Med* 2004; 37(12): 1951-62.
4. Kudin AP, Kudina TA, Seyfried J, Vielhaber S, Beck H, Elger CE, et al. Seizure-dependent modulation of mitochondrial oxidative phosphorylation in rat hippocampus. *Eur J Neurosci* 2002; 15(7): 1105-14.
5. Sharma DR, Sunkaria A, Bal A, Bhutia YD, Vijayaraghavan R, Flora SJ, et al. Neurobehavioral impairments, generation of oxidative stress and release of pro-apoptotic factors after chronic exposure to sulphur mustard in mouse brain. *Toxicol Appl Pharmacol* 2009; 240(2): 208-18.
6. Mehla J, Reeta KH, Gupta P, Gupta YK. Protective effect of curcumin against seizures and cognitive impairment in a pentylenetetrazole-kindled epileptic rat model. *Life Sci* 2010; 87(19-22): 596-603.
7. Gupta YK, Briyal S. Protective effect of vineatrol against kainic acid induced seizures, oxidative stress and on the expression of heat shock proteins in rats. *Eur Neuropsychopharmacol* 2006; 16(2): 85-91.
8. Sharma M, Briyal S, Gupta YK. Effect of alpha lipoic acid, melatonin and trans

- resveratrol on intracerebroventricular streptozotocin induced spatial memory deficit in rats. *Indian J Physiol Pharmacol* 2005; 49(4): 395-402.
9. Hosseini M, Harandizadeh F, Niazmand S, Soukhtanloo M, Mahmoudabady M. Antioxidant effect of *Achillea wilhelmsii* extract on pentylenetetrazole (seizure model)-induced oxidative brain damage in Wistar rats. *Indian J Physiol Pharmacol* 2013; 57(4): 418-24.
 10. Rosche J, Uhlmann C, Froscher W. Cognitive deficits and psychiatric disorders in patients with new-onset epilepsy. *Fortschr Neurol Psychiatr* 2010; 78(1): 18-26.
 11. Nematy M, Kamgar M, Mohajeri SM, Tabatabaei Zadeh SA, Jomezadeh MR, Akbarieh HO, et al. The effect of hydroalcoholic extract of *Coriandrum sativum* on rat appetite. *Avicenna J Phytomed* 2013; 3(1): 91-7.
 12. Rakhshandeh H, Sadeghnia HR, Ghorbani A. Sleep-prolonging effect of *Coriandrum sativum* hydro-alcoholic extract in mice. *Nat Prod Res* 2012; 26(22): 2095-8.
 13. Eidi M, Eidi A, Saeidi A, Molanaei S, Sadeghipour A, Bahar M, et al. Effect of coriander seed (*Coriandrum sativum* L.) ethanol extract on insulin release from pancreatic beta cells in streptozotocin-induced diabetic rats. *Phytother Res* 2009; 23(3): 404-6.
 14. Dhanapakiam P, Joseph JM, Ramaswamy VK, Moorthi M, Kumar AS. The cholesterol lowering property of coriander seeds (*Coriandrum sativum*): mechanism of action. *J Environ Biol* 2008; 29(1): 53-6.
 15. Samojlik I, Lakic N, Mimica-Dukic N, Dakovic-Svajcer K, Bozin B. Antioxidant and hepatoprotective potential of essential oils of coriander (*Coriandrum sativum* L.) and caraway (*Carum carvi* L.) (Apiaceae). *J Agric Food Chem* 2010; 58(15): 8848-53.
 16. de Almeida Melo E, Bion FM, Filho JM, Guerra NB. In vivo antioxidant effect of aqueous and ethereal coriander (*Coriandrum sativum* L.) extracts. *European Journal of Lipid Science and Technology* 2003; 105(9): 483-7.
 17. Harsha SN, Anilakumar KR. In vitro free radical scavenging and DNA damage protective property of *Coriandrum sativum* L. leaves extract. *J Food Sci Technol* 2014; 51(8): 1533-9.
 18. Tang EL, Rajarajeswaran J, Fung SY, Kanthimathi MS. Antioxidant activity of *Coriandrum sativum* and protection against DNA damage and cancer cell migration. *BMC Complement Altern Med* 2013; 13: 347.
 19. Sultana S, Ripa FA, Hamid K. Comparative antioxidant activity study of some commonly used spices in Bangladesh. *Pak J Biol Sci* 2010; 13(7): 340-3.
 20. Wangenstein H, Samuelsen AB, Malterud KE. Antioxidant activity in extracts from coriander. *Food Chemistry* 2004; 88(2): 293-7.
 21. Rakhshandeh H, Hosseini M. Potentiation of pentobarbital hypnosis by *Rosa damascena* in mice. *Indian J Exp Biol* 2006; 44(11): 910-2.
 22. Ebrahimzadeh Bideskan AR, Hosseini M, Mohammadpour T, Karami R, Khodamoradi M, Nemati KH, et al. Effects of soy extract on pentylenetetrazol-induced seizures in ovariectomized rats. *Zhong Xi Yi Jie He Xue Bao* 2011; 9(6): 611-8.
 23. Hosseini M, Ghasemzadeh RM, Sadeghnia HR, Rakhshandeh H. Effects of different extracts of *Rosa damascena* on pentylenetetrazol-induced seizures in mice. *Zhong Xi Yi Jie He Xue Bao* 2011; 9(10): 1118-24.
 24. Hosseini M, Sadeghnia HR, Salehabadi S, Alavi H, Gorji A. The effect of L-arginine and L-NAME on pentylenetetrazole induced seizures in ovariectomized rats, an in vivo study. *Seizure* 2009; 18(10): 695-8.
 25. Hosseini M, Harandizadeh F, Niazmand S, Soukhtanloo M, Faizpour A, Ghasemabady M. The role for nitric oxide on the effects of hydroalcoholic extract of *Achillea wilhelmsii* on seizure. *Avicenna J Phytomed* 2014; 4(4): 251-9.
 26. Kamkar AM, Nazariبورun A, Hosseini M. Analgesic effect of the aqueous and ethanolic extracts of clove. *Avicenna J Phytomed* 2013; 3(2): 186-92.
 27. Hosseini M, Pkan P, Rakhshandeh H, Aghaie A, Sadeghnia HR, Ghasemzadeh Rahbardi M. The Effect of Hydro-Alcoholic Extract of Citrus Flower on Pentylenetetrazole and Maximal Electroshock-Induced Seizures in Mice. *World Applied Sciences Journal* 2011; 15(8): 1104-9.
 28. Hosseini M, Pourganji M, Khodabandehloo F, Soukhtanloo M, Zabihi H. Protective Effect of L-Arginine against Oxidative Damage as a Possible Mechanism of its Beneficial Properties on Spatial Learning in Ovariectomized Rats. *Basic & Clinical Neuroscience* 2012; 3(5): 36-44.
 29. Vafae F, Hosseini M, Sadeghnia HR, Hadjzadeh MA, Soukhtanloo M, Rahimi M. The effects of soy extract on spatial learning and memory damage induced by global ischemia in ovariectomized rats. *Malays J Med Sci* 2014; 21(3): 19-30.
 30. Pourganji M, Hosseini M, Soukhtanloo M, Zabihi H, Hadjzadeh MA. Protective role of endogenous ovarian hormones against learning and memory impairments and brain tissues oxidative damage induced by lipopolysaccharide. *Iran Red Crescent Med J* 2014; 16(3): e13954.
 31. Costello DJ, Delanty N. Oxidative injury in epilepsy: potential for antioxidant therapy? *Expert Rev Neurother* 2004; 4(3): 541-53.
 32. Zhen J, Qu Z, Fang H, Fu L, Wu Y, Wang H, et al. Effects of grape seed proanthocyanidin extract on pentylenetetrazole-induced kindling and associated cognitive impairment in rats. *Int J Mol Med* 2014; 34(2): 391-8.
 33. Sejima H, Ito M, Kishi K, Noda A, Serikawa T. Regional excitatory and inhibitory amino acid concentrations in Noda epileptic rat (NER) brain. *Brain Dev* 1999; 21(6): 382-5.
 34. Porter RJ, Cereghino JJ, Gladding GD, Hessie BJ, Kupferberg HJ, Scoville B, et al. Antiepileptic Drug Development Program. *Cleve Clin Q* 1984; 51(2): 293-305.
 35. Hosseinzadeh H, Sadeghnia HR. Protective effect of safranal on pentylenetetrazol-induced seizures in the rat: involvement of GABAergic and opioids systems. *Phytomedicine* 2007; 14(4): 256-62.
 36. Loscher W, Honack D, Fassbender CP, Nolting B. The role of technical, biological and pharmacological factors in the laboratory evaluation of anticonvulsant drugs. III. Pentylenetetrazole seizure models. *Epilepsy Res* 1991; 8(3): 171-89.
 37. Xie T, Wang WP, Mao ZF, Qu ZZ, Luan SQ, Jia LJ, et al. Effects of epigallocatechin-3-gallate on pentylenetetrazole-induced kindling, cognitive impairment and oxidative stress in rats. *Neurosci Lett* 2012; 516(2): 237-41.
 38. Liu SH, Chang CD, Chen PH, Su JR, Chen CC, Chung HC. Docosahexaenoic acid and phosphatidylserine supplementations improve antioxidant activities and cognitive functions of the developing brain on pentylenetetrazol-induced seizure model. *Brain Res* 2012; 1451: 19-26.
 39. Sudha K, Rao AV, Rao A. Oxidative stress and antioxidants in epilepsy. *Clin Chim Acta* 2001; 303(1-2): 19-24.
 40. Rodrigues AD, Scheffel TB, Scola G, Santos MT, Fank B, de Freitas SC, et al. Neuroprotective and anticonvulsant effects of organic and conventional purple grape juices on seizures in Wistar rats induced by pentylenetetrazole. *Neurochem Int* 2012; 60(8): 799-805.
 41. Reilly C, Agnew R, Neville BG. Depression and anxiety in childhood epilepsy: a review. *Seizure* 2011; 20(8): 589-97.
 42. Maldonado A, Ramos W, Perez J, Huaman LA, Gutierrez EL. Convulsive status epilepticus: clinico-epidemiologic characteristics and risk factors in Peru. *Neurologia* 2010; 25(8): 478-84.
 43. Liang LP, Beaudoin ME, Fritz MJ, Fulton R, Patel M. Kainate-induced seizures, oxidative stress and neuronal loss in aging rats. *Neuroscience* 2007; 147(4): 1114-8.
 44. Golechha M, Bhatia J, Arya DS. Hydroalcoholic extract of *Embilca officinalis* Gaertn. affords protection against PTZ-induced seizures, oxidative stress and cognitive impairment in rats. *Indian J Exp Biol* 2010; 48(5): 474-8.
 45. Velaga MK, Yallapragada PR, Williams D, Rajanna S, Bettaiya R. Hydroalcoholic seed extract of *Coriandrum sativum* (Coriander) alleviates lead-induced oxidative stress in different regions of rat brain. *Biol Trace Elem Res* 2014; 159(1-3): 351-63.
 46. Cioanca O, Hritcu L, Mihasan M, Trifan A, Hancianu M. Inhalation of coriander volatile oil increased anxiolytic-antidepressant-like behaviors and decreased oxidative status in beta-amyloid (1-42) rat model of Alzheimer's disease. *Physiol Behav* 2014; 131: 68-74.
 47. Cioanca O, Hritcu L, Mihasan M, Hancianu M. Cognitive-enhancing and antioxidant activities of inhaled coriander volatile oil in amyloid beta(1-42) rat model of Alzheimer's disease. *Physiol Behav* 2013; 120: 193-202.
 48. Shila S, Kokilavani V, Subathra M, Panneerselvam C. Brain regional responses in antioxidant system to alpha-lipoic acid in arsenic intoxicated rat. *Toxicology* 2005; 210(1): 25-36.
 49. Devi SA, Kiran TR. Regional responses in antioxidant system to exercise training and dietary vitamin E in aging rat brain. *Neurobiol Aging* 2004; 25(4): 501-8.
 50. Soszynski M, Bartosz G. Decrease in accessible thiols as an index of oxidative damage to membrane proteins. *Free Radic Biol Med* 1997; 23(3): 463-9.
 51. Cardenas-Rodriguez N, Coballase-Urrutia E, Perez-Cruz C, Montesinos-Correa H, Rivera-Espinosa L, Sampieri A, III, et al. Relevance of the Glutathione System in Temporal Lobe Epilepsy: Evidence in Human and

- Experimental Models. *Oxid Med Cell Longev* 2014; 2014: 759293.
52. Anderson ME. Glutathione: an overview of biosynthesis and modulation. *Chem Biol Interact* 1998; 111-112: 1-14.
 53. Chen TS, Richie JP, Lang CA. The effect of aging on glutathione and cysteine levels in different regions of the mouse brain. *Proc Soc Exp Biol Med* 1989; 190(4): 399-402.
 54. Abbott LC, Nejad HH, Bottje WG, Hassan AS. Glutathione levels in specific brain regions of genetically epileptic (tg/tg) mice. *Brain Res Bull* 1990; 25(4): 629-31.
 55. Lal AA, Kumar T, Murthy PB, Pillai KS. Hypolipidemic effect of *Coriandrum sativum* L. in triton-induced hyperlipidemic rats. *Indian J Exp Biol* 2004; 42(9): 909-12.
 56. Hosseinzadeh H, Madanifard M. Anticonvulsant effects of coriandrum sativum L. Seed extracts in mice. *Archives of Iranian Medicine* 2000; 3(4): 81-4.
 57. Emamghoreishi M, Heidari-Hamedani G. Anticonvulsant effect of extract and essential oil of *Coriandrum sativum* seed in concious mice. *Iran J Pharm Res* 2004; 3(1): 71.
 58. Kunzemann J, Herrmann K. Isolation and identification of flavon(ol)-O-glycosides in caraway (*Carum carvi* L.), fennel (*Foeniculum vulgare* Mill.), anise (*Pimpinella anisum* L.), and coriander (*Coriandrum sativum* L.), and of flavon-C-glycosides in anise. I. Phenolics of spices (author's transl). *Z Lebensm Unters Forsch* 1977; 164(3): 194-200.
 59. Nassiri-Asl M, Mortazavi SR, Samiee-Rad F, Zangivand AA, Safdari F, Saroukhani S, et al. The effects of rutin on the development of pentylentetrazole kindling and memory retrieval in rats. *Epilepsy Behav* 2010; 18(1-2): 50-3.
 60. Sefil F, Kahraman I, Dokuyucu R, Gokce H, Ozturk A, Tutuk O, et al. Ameliorating effect of quercetin on acute pentylentetrazole induced seizures in rats. *Int J Clin Exp Med* 2014; 7(9): 2471-7.
 61. Medina JH, Viola H, Wolfman C, Marder M, Wasowski C, Calvo D, et al. Overview--flavonoids: a new family of benzodiazepine receptor ligands. *Neurochem Res* 1997; 22(4): 419-25.
 62. Griebel G, Perrault G, Tan S, Schoemaker H, Sanger DJ. Pharmacological studies on synthetic flavonoids: comparison with diazepam. *Neuropharmacology* 1999; 38(7): 965-77.
 63. Paladini AC, Marder M, Viola H, Wolfman C, Wasowski C, Medina JH. Flavonoids and the central nervous system: from forgotten factors to potent anxiolytic compounds. *J Pharm Pharmacol* 1999; 51(5): 519-26.
 64. Youdim KA, Shukitt-Hale B, Joseph JA. Flavonoids and the brain: interactions at the blood-brain barrier and their physiological effects on the central nervous system. *Free Radic Biol Med* 2004; 37(11): 1683-93.
 65. Kang TH, Jeong SJ, Kim NY, Higuchi R, Kim YC. Sedative activity of two flavonol glycosides isolated from the flowers of *Albizia julibrissin* Durazz. *J Ethnopharmacol* 2000; 71(1-2): 321-3.
 66. Elisabetsky E, Brum LF, Souza DO. Anticonvulsant properties of linalool in glutamate-related seizure models. *Phytomedicine* 1999; 6(2): 107-13.
 67. de Sousa DP, Nobrega FF, Santos CC, de Almeida RN. Anticonvulsant activity of the linalool enantiomers and racemate: investigation of chiral influence. *Nat Prod Commun* 2010; 5(12): 1847-51.
 68. Usta J, Kreydiyyeh S, Knio K, Barnabe P, Bou-Moughlabay Y, Dagher S. Linalool decreases HepG2 viability by inhibiting mitochondrial complexes I and II, increasing reactive oxygen species and decreasing ATP and GSH levels. *Chem Biol Interact* 2009; 180(1): 39-46.