

- Mol Med 2009; 41(1): 33-41.
18. McDonald WI, Compston A, Edan G, Goodkin D, Hartung HP, Lublin FD, et al. Recommended diagnostic criteria for multiple sclerosis: Guidelines from the International Panel on the diagnosis of multiple sclerosis. *Ann Neurol* 2001; 50(1): 121-7.
 19. McPherson MJ, Moller SG, Howe BC. PCR: The Basics from Background to Bench. New York, NY: Springer-Verlag; 2000. p. 23-119.
 20. Compston A, Coles A. Multiple sclerosis. *Lancet* 2008; 372(9648): 1502-17.
 21. Beecham AH, Patsopoulos NA, Xifara DK, Davis MF, Kempainen A, Cotsapas C, et al. Analysis of immune-related loci identifies 48 new susceptibility variants for multiple sclerosis. *Nat Genet* 2013; 45(11): 1353-60.
 22. Pearce EL, Mullen AC, Martins GA, Krawczyk CM, Hutchins AS, Zediak VP, et al. Control of effector CD8+ T cell function by the transcription factor Eomesodermin. *Science* 2003; 302(5647): 1041-3.
 23. Sawcer S, Hellenthal G, Pirinen M, Spencer CC, Patsopoulos NA, Moutsianas L, et al. Genetic risk and a primary role for cell-mediated immune mechanisms in multiple sclerosis. *Nature* 2011; 476(7359): 214-9.
 24. Maurano MT, Humbert R, Rynes E, Thurman RE, Haugen E, Wang H, et al. Systematic localization of common disease-associated variation in regulatory DNA. *Science* 2012; 337(6099): 1190-5.
 25. Hindorf LA, Sethupathy P, Junkins HA, Ramos EM, Mehta JP, Collins FS, et al. Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. *Proc Natl Acad Sci U S A* 2009; 106(23): 9362-7.
 26. O'Connor RA, Cambrook H, Huettnner K, Anderton SM. T-bet is essential for Th1-mediated, but not Th17-mediated, CNS autoimmune disease. *Eur J Immunol* 2013; 43(11): 2818-23.
 27. Bettelli E, Sullivan B, Szabo SJ, Sobel RA, Glimcher LH, Kuchroo VK. Loss of T-bet, but not STAT1, prevents the development of experimental autoimmune encephalomyelitis. *J Exp Med* 2004; 200(1): 79-87.
 28. Hirota K, Duarte JH, Veldhoen M, Hornsby E, Li Y, Cua DJ, et al. Fate mapping of IL-17-producing T cells in inflammatory responses. *Nat Immunol* 2011; 12(3): 255-63.
 29. Chan RW, Lai FM, Li EK, Tam LS, Chow KM, Li PK, et al. Expression of T-bet, a type 1 T-helper cell transcription factor, in the urinary sediment of lupus patients predicts disease flare. *Rheumatology (Oxford)* 2007; 46(1): 44-8.
 30. Fyall KM, Fong AM, Rao SB, Ibrahim IC, Waxweiler WT, Thomas NE, et al. TBX21 transcription factor T-1514C polymorphism is associated with decreased IFN-gamma and IL-4 production by primary human lymphocytes. *Hum Immunol* 2012; 73(6): 673-6.
 31. Drulovic J, Savic E, Pekmezovic T, Mesaros S, Stojisavljevic N, Dujmovic-Basuroski I, et al. Expression of Th1 and Th17 cytokines and transcription factors in multiple sclerosis patients: does baseline T-bet mRNA predict the response to interferon-beta treatment? *J Neuroimmunol* 2009; 215(1-2): 90-5.
 32. Li JR, Li JG, Deng GH, Zhao WL, Dan YJ, Wang YM, et al. A common promoter variant of TBX21 is associated with allele specific binding to Yin-Yang 1 and reduced gene expression. *Scand J Immunol* 2011; 73(5): 449-58.
 33. Chen S, Zhao W, Tan W, Luo X, Dan Y, You Z, et al. Association of TBX21 promoter polymorphisms with type 1 autoimmune hepatitis in a Chinese population. *Hum Immunol* 2011; 72(1): 69-73.
 34. Veeger C, Brodbeck K, Sawadogo M, Haring HU, Schleicher ED. Upstream regulatory factor (USF) proteins induce human TGF-beta1 gene activation via the glucose-response element-1013/-1002 in mesangial cells: up-regulation of USF activity by the hexosamine biosynthetic pathway. *J Biol Chem* 2004; 279(16): 15908-15.
 35. Pender MP, Greer JM. Immunology of multiple sclerosis. *Curr Allergy Asthma Rep* 2007; 7(4): 285-92.

Proof VOSIORS