

East Asia and demonstrated the favorable use of DMF in patients with RRMS, which was similar to the findings of our study. In their study, DMF could successfully reduce new GD lesions from baseline to week 24 by 75% and reduce the number of relapses by 42% over a 24-week period.²²

Conclusion

Relapsing MS is a chronic neurologic disorder that can be managed without significant complications if appropriate treatment is identified. DMF is a newly-used drug for treatment of relapsing MS and is still being studied for long-term effects in various clinical trials. The results of the present clinical trial showed that treatment of relapsing MS with DMF would provide

decreased attack rate and GD lesions. Moreover, most of the patients were satisfied with their treatment and EDSS score was significantly reduced without any serious complications, which may result in termination of DMF therapy.

Conflict of Interests

The authors declare no conflict of interest in this study.

Acknowledgments

We would like to express our sincere gratitude and appreciation to School of Medicine of Mashhad University of Medical Sciences as well as Mashhad Comprehensive MS Center for their valuable cooperation throughout the course of the project.

References

- Browne P, Chandraratna D, Angood C, Tremlett H, Baker C, Taylor BV, et al. Atlas of Multiple Sclerosis 2013: A growing global problem with widespread inequity. *Neurology* 2014; 83(11): 1022-4.
- Mansouri S, Zayeri F. Global and regional trends of multiple sclerosis disability-adjusted life years rates: A 25-year assessment. *Neuroepidemiology* 2019; 52(1-2): 17-24.
- Thompson AJ, Banwell BL, Barkhof F, Carroll WM, Coetzee T, Comi G, et al. Diagnosis of multiple sclerosis: 2017 revisions of the McDonald criteria. *Lancet Neurol* 2018; 17(2): 162-73.
- Gold R, Kappos L, Arnold DL, Bar-Or A, Giovannoni G, Selmaj K, et al. Placebo-controlled phase 3 study of oral BG-12 for relapsing multiple sclerosis. *N Engl J Med* 2012; 367(12): 1098-107.
- Fox RJ, Miller DH, Phillips JT, Hutchinson M, Havrdova E, Umeta M, et al. Placebo-controlled phase 3 study of oral BG-12 or glatiramer in multiple sclerosis. *N Engl J Med* 2012; 366(12): 1001-97.
- Setayeshgar S, Khatami E, Zhu F, Zhang X, Zhang T, Marrie RA, et al. Use of the new oral disease-modifying therapies for multiple sclerosis in British Columbia, Canada: The first five-years. *Mult Scler Relat Dis* 2018; 25: 57-60.
- Braune S, Grimm S, van Hovell P, Freudensprung U, Pellegrini F, Hyde R, et al. Comparative effectiveness of delayed-release dimethyl fumarate versus interferon, glatiramer acetate, teriflunomide, or fingolimod: Results from the German NeuroTransData registry. *J Neurol* 2018; 265(12): 2980-92.
- Ontaneda D, Nicholas J, Carraro M, Zhou J, Hou Q, Babb J, et al. Comparative effectiveness of dimethyl fumarate versus fingolimod and teriflunomide among MS patients switching from first-generation platform therapies in the US. *Mult Scler Relat Disord* 2019; 27: 101-11.
- Conde S, Moisset X, Pereira B, Zuel M, Colamarino R, Maillet-Vioud M, et al. Dimethyl fumarate and teriflunomide for multiple sclerosis in a real-life setting: A French retrospective cohort study. *Eur J Neurol* 2019; 26(3): 460-7.
- Mirabella M, Prosperini L, Lucchini M, Boffa L, Borriello G, Buscarinu MC, et al. Safety and efficacy of dimethyl fumarate in multiple sclerosis: An Italian, multicenter, real-world study. *CNS Drugs* 2018; 32(10): 963-70.
- Kresa-Rezaei K, Repovic P, Robertson D, Okwuonye N, Meizer L, Mendoza JP. Effectiveness of delayed-release dimethyl fumarate on clinical and patient-reported outcomes in patients with relapsing multiple sclerosis switching from glatiramer acetate: RESPOND, a prospective observational study. *Clin Ther* 2018; 40(12): 2077-87.
- Gold R, Giovannoni G, Phillips JT, Fox RJ, Zhang A, Meltzer L, et al. Efficacy and safety of delayed-release dimethyl fumarate in patients newly diagnosed with relapsing-remitting multiple sclerosis (RRMS). *Mult Scler* 2015; 21(1): 57-66.
- D'Amico E, Zanghi A, Callari G, Borriello G, Gallo A, Graziano G, et al. Comparable efficacy and safety of dimethyl fumarate and teriflunomide treatment in relapsing-remitting multiple sclerosis: An Italian real-world multicenter experience. *Ther Adv Neurol Disord* 2018; 11: 1756286418796404.
- Kalincik T, Kubala HE, Horakova D, Izquierdo G, Prat A, Girard M, et al. Comparison of fingolimod, dimethyl fumarate and teriflunomide for multiple sclerosis. *J Neurol Neurosurg Psychiatry* 2019; 90(4): 458-68.
- Mallucci G, Annovazzi P, Miente S, Torri-Clerici V, Matta M, La GS, et al. Two-year real-life efficacy, tolerability and safety of dimethyl fumarate in an Italian multicentre study. *J Neurol* 2018; 265(8): 1850-9.
- Properino L, Lucchini M, Haggiag S, Balantonio P, Bianco A, Buscarinu MC, et al. Fingolimod vs dimethyl fumarate in multiple sclerosis: A real-world propensity score-matched study. *Neurology* 2018; 91(2): e153-e161.
- Huisman E, Papadimitropoulou K, Jarrett J, Bending M, Firth Z, Allen F, et al. Systematic literature review and network meta-analysis in highly active relapsing-remitting multiple sclerosis and rapidly evolving severe multiple sclerosis. *BMJ Open* 2017; 7(3): e013430.
- Fogarty E, Schmitz S, Tubridy N, Walsh C, Barry M. Comparative efficacy of disease-modifying therapies for patients with relapsing remitting multiple sclerosis: Systematic review and network meta-analysis. *Mult Scler Relat Disord* 2016; 9: 23-30.
- Lebrun C, Rocher F. Cancer risk in patients with multiple sclerosis: Potential impact of disease-modifying drugs. *CNS Drugs* 2018; 32(10): 939-49.
- Kouliniska I, Riester K, Chalkias S, Edwards MR. Effect of bismuth subsalicylate on gastrointestinal tolerability in healthy volunteers receiving oral delayed-release dimethyl fumarate: PREVENT, a randomized, multicenter, double-blind, placebo-controlled study. *Clin Ther* 2018; 40(12): 2021-30.
- Ochi H, Niino M, Onizuka Y, Hiramatsu K, Hase M, Yun J, et al. 72-week safety and tolerability of dimethyl fumarate in Japanese patients with relapsing-remitting multiple sclerosis: Analysis of the randomised, double blind, placebo-controlled, phase III APEX study and its open-label extension. *Adv Ther* 2018; 35(10): 1598-611.
- Saida T, Yamamura T, Kondo T, Yun J, Yang M, Li J, et al. A randomized placebo-controlled trial of delayed-release dimethyl fumarate in patients with relapsing-remitting multiple sclerosis from East Asia and other countries. *BMC Neurol* 2019; 19(1): 5.