Current Journal of Neurology

Neurological Images

Curr J Neurol 2024; 23(3): 199



Ocular lateropulsion in acute vestibular neuritis

Received: 12 Mar. 2024 Accepted: 01 May 2024

Lisle Blackbourn, Maher Salem, Deepak Nair

Department of Neurology, College of Medicine, University of Illinois, Peoria, USA

Keywords

Stroke; Head Impulse Test; Saccades; Smooth Pursuit; Eye Movements; Nystagmus; Vestibular Neuritis

Patient was a 41-year-old man who presented with sudden-onset nausea, vertigo, and gait imbalance with a blood pressure on arrival of 230/132. On physical examination, the patient had left rotational nystagmus in upright position, ocular lateropulsion, and a positive horizontal head impulse test.

The ocular lateropulsion can be seen in video 1. Brain computed tomography (CT), head and neck CT angiography (CTA), and brain magnetic resonance imaging (MRI) all showed no acute abnormalities. Ocular lateropulsion can be seen in both strokes such as Wallenberg syndrome and in benign pathology such as acute vestibular neuritis.¹

The ocular lateropulsion usually also exhibits saccades in stroke versus a smooth pursuit in acute vestibular neuritis. A positive horizontal head impulse test usually points to a peripheral cause; however, it should be noted that patients with lateral pontine and cerebellar strokes can still have a positive horizontal head impulse test.² A negative head impulse test points to a central pathology.

How to cite this article: Blackbourn L, Salem M, Nair D. Ocular lateropulsion in acute vestibular neuritis. Curr J Neurol 2024; 23(3): 199.

Conflict of Interests

The authors declare no conflict of interest in this study.

Acknowledgments

None.

References

 Kattah JC, Badihian S, Pula JH, Tarnutzer AA, Newman-Toker DE, Zee DS. Ocular lateral deviation with brief removal of visual fixation differentiates central from peripheral

vestibular syndrome. J Neurol 2020; 267(12): 3763-72.
Newman-Toker DE, Kattah JC, Alvernia JE, Wang DZ. Normal head impulse test differentiates acute cerebellar strokes from vestibular neuritis. Neurology 2008; 70(24 Pt 2): 2378-85.

Copyright © 2024 Iranian Neurological Association, and Tehran University of Medical Sciences Published by Tehran University of Medical Sciences

Corresponding Author: Lisle Blackbourn Email: lblackbourn@uwalumni.com

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 international license (http://creativecommons.org/licenses/by-nc/4.0/). Non-commercial purposes uses of the work are permitted, provided the original work is properly cited.